

Elite Plastic Surgery

ROZBEH TORABI, MD ■ RADBEH TORABI, MD ■ KRISTA SINCLAIR, FNP-C, RNFA
Phoenix | 10910 N. TATUM BLVD, SUITE B-100, PHOENIX, AZ 85028 | Tel: (480) 291-6895 | Fax: (480) 948-3750
Chandler | 1120 S. Dobson Rd. Chandler Suite 110, AZ 85286 | P: (480) 937-0180 | Fax: (480) 948-3750

NECK LIFT

Post Operative Instructions

AT HOME AFTER NECK LIFT SURGERY:

- Always elevate your head. Avoid turning your head from side to side. To turn, move head and shoulders together. No lifting or bending over.
- Get out of bed with help. It is not good to lie still in bed, so move your feet and legs a couple of times every hour while you are awake.
- A liquid diet is best for the day of the surgery. Begin by taking liquids slowly and progress to soups or JELL-O®. You can start with a light diet the next day.
- You may have drains in place after surgery; if so, instructions will be given.
- You can expect some swelling of the face and eyes after surgery. Call if swelling on one side is more pronounced or harder than the other side; or if you are having pain that is not relieved by pain medication.
- Remember: DO NOT SMOKE or be around smoke. This is very important.
- While bandages are on, do not fall asleep in a chair.

OTHER POSTOPERATIVE NECK LIFT INSTRUCTIONS:

- Discourage visitors and keep facial movements (smiling, talking, chewing, yawning, etc.) to a minimum for the first few days.
- You will return to the office at post day 1 to have your drain removed.
- You will return to the office at post-op week 1 to have bandages and stitches removed.
- You may shower at post-op day 2 or 48 hours after your surgery.
- Wash your hair gently with baby shampoo and rinse thoroughly until the soap is out. It may take several washings before all the crusts are out of hair. Do not use hairsprays, conditioners, gels, etc. You may wash your face gently with a bland soap (Dove,

Elite Plastic Surgery

ROZBEH TORABI, MD ■ RADBEH TORABI, MD ■ KRISTA SINCLAIR, FNP-C, RNFA
Phoenix | 10910 N. TATUM BLVD, SUITE B-100, PHOENIX, AZ 85028 | Tel: (480) 291-6895 | Fax: (480) 948-3750
Chandler | 1120 S. Dobson Rd. Chandler Suite 110, AZ 85286 | P: (480) 937-0180 | Fax: (480) 948-3750

Neutrogena, etc.). You may use a moisturizer on your face, being careful to keep it away from the stitch lines.

- Makeup may be applied immediately, but do not get it near the stitch lines until two days after the stitches have been removed.
- Normal activities can be gradually resumed after two weeks. You may resume exercise after four-six weeks.
- The skin of your face is sensitive to sunlight after surgery. Protect your facial skin with sunscreen (SPF 15 or greater). Sauna and steam baths should be avoided for four weeks following surgery.

GENERAL INFORMATION:

- Swelling and bruising will almost always be more than you expected. Bruising generally subsides in seven days. Most of the swelling subsides in two weeks. It will continue to improve for two to three months.
- The scars have a tendency to redden and become more noticeable a few weeks after the surgery. However, the redness will fade with time.
- Numbness of the scalp is common. Often this begins to return in six weeks but may take six to nine months to improve.
- Discoloration of the whites of the eyes may occur if there's a lot of swelling. This is painless, will not harm your vision and will disappear completely.
- Arnica reduces swelling and bruising and can be found in natural food/nutrition stores. Begin taking two days before surgery.
- The office will contact you the day after surgery to check-up on you, answer questions and book follow-up. (Friday/Weekend surgeries - patients will be called on Monday.)

Elite Plastic Surgery

ROZBEH TORABI, MD ■ RADBEH TORABI, MD ■ KRISTA SINCLAIR, FNP-C, RNFA
Phoenix | 10910 N. TATUM BLVD, SUITE B-100, PHOENIX, AZ 85028 | Tel: (480) 291-6895 | Fax: (480) 948-3750
Chandler | 1120 S. Dobson Rd. Chandler Suite 110, AZ 85286 | P: (480) 937-0180 | Fax: (480) 948-3750

Medications Prescribed (not all medication may apply)

Medications Prescribed (not all medication may apply)

Please take all your medications as prescribed.

Scheduled Medications: Please take these medications as scheduled

Antibiotic: Keflex (cephalexin) 500mg (started day of surgery, is a SCHEDULED medication, please follow the prescription ordered) is an antibiotic you must take for a total of 5 days. (Note that antibiotics may impair birth control pills.)

(*May not apply to you) Blood thinner: Lovenox 30 mg/0.3 ml injectable solution (started day AFTER surgery) is used for the prevention of blood clots after surgery. You or your caregiver will administer 1 subcutaneous injection into your thighs daily for 5 days.

Tylenol (Acetaminophen) (Age GREATER than 60: 1000 mg every 8 hours) (Age LESS than 60: 1000 mg every 6 hours)

As needed: These medications are scheduled “as needed” per your pain level. We provide you with many types of medications that focus on the “type” of pain you are experiencing. If you plan to take more than one of these medications, please plan to take them at least 2 hours apart as they can make you sleepy.

Toradol (ketorolac): 10 mg every 6 hrs = 15 tabs

Type of Pain relief: Anti-inflammatory: Swelling/Pain: (started day of surgery as needed, please follow prescription ordered) This is a different form of Ibuprofen so please stay away from other anti-inflammatories while taking this medication.

Gabapentin (Neurontin): 100 mg; every 8 hours = 48 tabs (16 days)

Type of Pain relief: Nerve Pain: Itching/Burning/Tingling/Numbing/Sharp: (started day of surgery as needed, please follow the prescription ordered) is used for pain relief every 12 hours. Please let us know if you are taking any prescriptions for mood-altering (depression or bipolar) medications as these can potentially interact.

***this medication can make you sleepy, please do not plan to drive on this medication**

Cyclobenzaprine (Flexeril): 5 mg; every 8 hrs = 10 tabs

Type of Pain relief: Muscle Relaxer: Muscle spasms/Muscle cramping: (started day of surgery as needed, please follow prescription ordered) is a muscle spasm medication.

Elite Plastic Surgery

ROZBEH TORABI, MD ▫ RADBEH TORABI, MD ▫ KRISTA SINCLAIR, FNP-C, RNFA
Phoenix | 10910 N. TATUM BLVD, SUITE B-100, PHOENIX, AZ 85028 | Tel: (480) 291-6895 | Fax: (480) 948-3750
Chandler | 1120 S. Dobson Rd. Chandler Suite 110, AZ 85286 | P: (480) 937-0180 | Fax: (480) 948-3750

***please do not drive while on this medication**

Oxycodone (Narcotic Pain Reliever): po every 4-6 hours as needed = 30 tabs

Type of Pain Relief: General pain (started day of surgery as needed, please follow the prescription ordered) is an oral narcotic prescribed to manage your pain.

***please do not drive while on this medication**

Docusate Sodium (Colace): Stool Softener: every 12 hours = 14 tabs (7 days)

Senna: Bowel Stimulant: every 12 hours = 14 tabs (7 days)

You may take the Colace and Senna together.